

A strong dissipative inflationary model

Mauricio Bellini*

Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)

and

*Departamento de Física, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata,
Funes 3350, (7600) Mar del Plata, Buenos Aires, Argentina.*

The dynamics of a single scalar field which describes a second order phase transition is considered taking into account the thermal effects in the high temperature limit. I conclude that thermal effects play a very important role during inflation and do not must to be ignored in models for the early universe. At the end of inflation thermal fluctuations are more important than quantum fluctuations, due to the fact that $T > H_c$.

Pacs numbers: 98.80.Cq

I. INTRODUCTION

An inflationary phase eliminates the major problems in cosmology, namely flatness, the horizon problem, homogeneity and the numerical density of monopoles [1–6]. However, the main problem is how to attach the observed universe to the end of the inflationary epoch. In the standard (isentropic) inflationary scenarios, the radiation energy density ρ_r scales with the inverse fourth power of the scale factor, becoming quickly negligible. In such case, a short time reheating period terminates the inflationary period and initiates the radiation dominated epoch. On the other hand, the only condition required by general relativity for inflation is that $\rho_r < \rho_v$. Inflation in presence of non-negligible radiation is characterized by nonisentropic expansion, and thermal seeds of density fluctuations. Within a realistic quantum field theory model, Hosoya and Sakagami [7] initially formulated dissipative dynamics. In their formulation, the near-thermal-equilibrium dynamics is expressed through an expansion involving equilibrium correlation functions. Subsequently Morikawa [8] formulated the same problem in terms of an elegant real time finite temperature field theory formalism. These works were developed further by Lawrie [9] and Gleiser and Ramos [10]. For models like warm inflation [11] the overdamped regime of the inflaton is the one of interest. A realization of overdamped motion within quantum field theory model based on this formulation was obtained in [12]. A more recently warm inflationary model inspired in string theories was developed in [13].

In this paper I study the equation of motion for the expectation value of the inflaton field φ on a thermodynamical average for a symmetric massive scalar field with self-interaction. Unfortunately no reliable theory of a system far from equilibrium has yet been available. We have to content ourselves to work in the hydrodynamical regime, i.e., slight-off-equilibrium state following some ideas of Hosoya and Sakagami [7], who obtained this equation which also includes the effect of the temperature increase by friction during inflation. For this propose I will consider the equation of motion $\square\varphi + V'(\varphi) = 0$ on a background spatially flat Friedmann-Robertson-Walker (FRW) $ds^2 = -dt^2 + a^2dx^2$ metric on a thermodynamical average by means of the semiclassical approach $\varphi(\vec{x}, t) = \phi_c(t) + \phi(\vec{x}, t)$. Here, $\phi_c(t)$ is given by $\langle\varphi(\vec{x}, t)\rangle = \text{tr}(\rho\varphi)$ for a suitable nonequilibrium statistical operator ρ and $\phi(\vec{x}, t)$ is the fluctuation field around the thermodynamical average value ϕ_c . Furthermore, the operator ϕ has a zero thermodynamical average $\langle\phi\rangle = 0$. In the Heisenberg representation the operator ρ is time independent (i.e., $\frac{d\rho}{dt} = 0$). The classical scalar field ϕ_c starts at a point a little bit away from the top and then rolls down along the long slope to the minimum of the effective potential. If the friction force is strong enough, there will be no oscillation and ϕ_c simply slowly rolls down to the bottom.

II. THE EFFECTIVE EQUATION OF MOTION

We consider the thermodynamical average of the field equation $\square\varphi + V'(\varphi) = 0$

$$\langle\square\varphi + V'(\varphi)\rangle = 0, \quad (1)$$

*E-mail address: mbellini@mdp.edu.ar

where

$$\langle \square \varphi \rangle = (\ddot{\phi}_c + \ddot{\phi}) - \frac{1}{a^2} \nabla^2 \phi + 3 \left\langle H(\phi_c + \phi)(\dot{\phi}_c + \dot{\phi}) \right\rangle.$$

For simplicity, I will consider $\langle H(\phi_c + \phi) \rangle \simeq H_c(\phi_c)$ where $H_c(\phi_c) = \dot{a}/a$ is the Hubble parameter and a is the scale factor of the universe. The dot and prime denote the derivatives with respect to time and the field, respectively. On the other hand, the thermodynamical average $\langle V'(\varphi) \rangle$ can be expanded as

$$\langle V'(\phi_c + \phi) \rangle = V'(\phi_c) + V''(\phi_c) \langle \phi \rangle + \frac{V'''(\phi_c)}{2!} \langle \phi^2 \rangle + \dots \quad (2)$$

Hence, the equation of motion for the classical field being given by

$$\ddot{\phi}_c + 3H_c \dot{\phi}_c + V'(\phi_c) + \frac{V'''(\phi_c)}{2!} \langle \phi^2 \rangle + \dots = 0. \quad (3)$$

The thermodynamical average of the squared fluctuations are

$$\langle \phi^2 \rangle = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\omega(p)} n(p), \quad (4)$$

where $\omega^2(p) = [\vec{p}^2 + V''(\phi_c)]$ and $n(p)$ is the bosons distribution given by $n(p) = n_0(p) + \delta n(p)$ in the first order approximation. The local equilibrium distribution $n_0(p)$ is given by

$$n_0(p) = \frac{1}{e^{\beta \omega(p)} - 1}, \quad (5)$$

where $\beta = 1/T$ is the inverse of the temperature T . Furthermore, $\delta n(p)$ is a small deviation of n_0 . If τ is the relaxation time of the fluctuations, in the relaxation-time approximation $\delta n = -\tau \partial_t n_0$

$$\delta n = \tau n_0 \left[\frac{\beta}{2\omega(p)} V'''(\phi_c) \dot{\phi}_c + \dot{\beta} \omega \right], \quad (6)$$

where was used $\dot{\omega} = \frac{V'''(\phi_c) \dot{\phi}_c}{2\omega}$ and $d_t \equiv \dot{\phi}_c d_{\phi_c}$. Furthermore, τ is the relaxation time for the fluctuations.

A. Calculation of $\langle \phi^2 \rangle$

From eqs. (4), (5) and (6) the thermodynamical average of the squared fluctuations ϕ are

$$\begin{aligned} \langle \phi^2 \rangle &= \langle \phi^2 \rangle_0 + \left[\dot{\phi}_c V'''(\phi_c) \int \frac{d^3 p}{(2\pi)^3} \frac{\tau n_0(1 + n_0)\beta}{2\omega^2(p)} \right. \\ &\quad \left. + \int \frac{d^3 p}{(2\pi)^3} \tau n_0(1 + n_0) \dot{\beta} \right], \end{aligned} \quad (7)$$

where the first term in eq. (7) gives the fluctuations in thermal equilibrium

$$\langle \phi^2 \rangle_0 = \int \frac{d^3 p}{(2\pi)^3} \frac{n_0(p)}{\omega(p)}, \quad (8)$$

and the remaining terms in eq. (7) describe the contribution out of the thermal equilibrium. Replacing eq. (5) in the equation (8) and integrating, we obtain the following result in the high temperature limit (i.e., for $\beta m \ll 1$):

$$\langle \phi^2 \rangle_0 \simeq \frac{T^2}{12} \left(1 - \frac{4m}{\pi^2 T} \right). \quad (9)$$

On the other hand, using the fact that $\tau^{-1} \simeq \frac{\lambda^4}{3.2^9 \pi} \frac{T^2}{\omega}$ [7] in the high temperature limit, the first integrate in eq. (7) gives

$$\int \frac{d^3p}{(2\pi)^3} \frac{\tau n_0(1+n_0)\beta}{\omega^2(p)} \beta \simeq \frac{3.2^8}{\pi T} \ln(T/m). \quad (10)$$

The second integrate in eq. (7) results

$$\int \frac{d^3p}{(2\pi)^3} \tau n_0(1+n_0) \dot{\beta} \simeq -\frac{3.2^7}{\pi} \dot{T}. \quad (11)$$

We can obtain $\dot{T}$ from the equation which describes the evolution for $\rho_r = \frac{\pi^2}{30} g_{eff} T^4$

$$\dot{\rho}_r + 4H_c \rho_r - \delta = 0. \quad (12)$$

If there were no dissipation, then $\delta = 0$ and the radiation component would be rapidly red-shifted away as $\rho_r \sim e^{-int4H_c dt}$. However with dissipation, radiation is being produced continuously from conversion of scalar field vacuum energy. In our case $\delta = \eta \dot{\phi}_c^2$ describes the interaction term due to the decay of the inflaton field without consider the temporal evolution of the temperature due to the expansion of the universe. We are interested in the study of models where $\dot{\rho}_r > 0$ (Fresh inflation) [14]. Inflationary models where the radiation energy density remains almost constant ($\dot{\rho}_r \simeq 0$) are named Warm Inflation [11]. I will consider the potential

$$V(\phi_c) = \frac{m^2}{2} \phi_c^2 + \frac{\lambda^2}{24} \phi_c^4 + \Lambda, \quad (13)$$

where Λ is the cosmological constant. For this potential we obtain the following dissipative parameter:

$$\begin{aligned} \eta &= \beta \left[\frac{V'''(\phi_c)}{2} \right]^2 \int \frac{d^3p}{(2\pi)^3} \tau \frac{n_0}{\omega^2} (1+n_0) \\ &\simeq \frac{3.2^6}{\pi} \frac{\phi_c^2}{T} \ln\left(\frac{T}{m}\right). \end{aligned} \quad (14)$$

Hence, from eqs. (12) and (14), we obtain the temporal evolution for T in eq. (11)

$$\dot{T} = \frac{45.2^5}{\pi^3 g_{eff}} \frac{\ln(T/m)}{T^4} \dot{\phi}_c^2 \phi_c^2 - H_c T. \quad (15)$$

Note that eq. (15) describes the evolution of the temperature taking into account dissipation and expansion effects. The first term in eq. (15) comes from dissipative effects and the second one takes into account the expansion of the universe. For this reason, for $\eta = 0$ we obtain the red-shifted temperature $T \sim e^{-\int H_c dt}$ (which would be $T \sim e^{-H_0 t}$, with $H_0 = \text{const.}$, in a de Sitter expansion). In our case dissipative effects are more important than the expansion of the universe (i.e., $\eta > H_c$), so that the temperature increases: $\dot{T} > 0$.

Finally, the squared fluctuations $\langle \phi^2 \rangle$ in the high temperature limit, are

$$\begin{aligned} \langle \phi^2(\vec{x}, t) \rangle|_{T \gg m} &\simeq \frac{T^2}{12} \left[1 - \frac{4m}{\pi^2 T} \right] \\ &+ \frac{3.2^7}{\pi} \left\{ \frac{\ln(T/m)}{T} \left[\dot{\phi}_c \phi_c - \frac{45.2^5}{\pi^3 g_{eff} T^3} \phi_c^2 \dot{\phi}_c^2 \right] + H_c T \right\}. \end{aligned} \quad (16)$$

B. Inflationary dynamics

Once obtained the equation that describes the temporal evolution for the temperature [see eq. (15)] and the squared field fluctuations $\langle \phi^2 \rangle$ we can write the effective equation of motion for the background field ϕ_c

$$\ddot{\phi}_c + [3H_c + \eta_{eff}] \dot{\phi}_c + V'_{eff}(\phi_c, T) = 0, \quad (17)$$

with η_{eff} and V'_{eff} given by

$$\eta_{eff} = \eta - \frac{135.2^{11} \ln(T/m)}{\pi^4 g_{eff}} \frac{\phi_c^3 \dot{\phi}_c}{T^4}, \quad (18)$$

$$V'_{eff}(\phi_c, T) = \left[m^2 + \frac{\lambda^2 T^2}{24} \left(1 - \frac{4m}{\pi^2 T} \right) \right] \phi_c + \frac{\lambda}{6} \phi_c^3 + \frac{3.2^6}{\pi} H_c \phi_c T. \quad (19)$$

The additional term in η_{eff} comes from the first term in eq. (15) which is originated by the interaction of the inflaton field with the background thermal bath. In other words η_{eff} is given by the η -self-interaction of the inflaton field plus the additional dissipative contribution given by the fact that the fields produced by self-interaction interact with the fields of the background thermal bath. Furthermore, the remaining term in eq. (15) is conservative and can be added in the ϕ_c -derivative of the effective potential $V_{eff}(\phi_c, T)$ [see eq. (19)]. This term is originated by the expansion of the universe and its origin is purely cosmological.

It is very important to take into account that the equation (17) is valid only if the hydrodynamical condition is satisfied:

$$\tau \ll \left| \frac{a}{\dot{a}} \right|, \left| \frac{\phi_c}{\dot{\phi}_c} \right|.$$

This means that the relaxation time of the fluctuations must to be very small with respect to the inverse of the Hubble parameter and the characteristic time of change for the background field ϕ_c . Furthermore, one has to be careful about the region in which eq. (17) is applicable. This equation is only valid in the limit of high temperatures and the condition $T/m \gg 1$ must to be fulfilled.

Finally, the Friedmann equation $\langle H^2(\varphi) \rangle = 8\pi G/3 [\langle \rho_\phi + \rho_r \rangle] \simeq 8\pi G/3 [\langle \dot{\phi}^2/2 \rangle + \langle V(\varphi) \rangle]$ for an inflating universe with potential (13) that expands in a background thermal bath can be written as [see eq. (19)]

$$H_c^2(\phi_c) + H_c(\phi_c) H_c''(\phi_c) \langle \phi^2 \rangle = \frac{8\pi G}{3} \left[\frac{\dot{\phi}_c^2}{2} + V(\phi_c) + \frac{V''(\phi_c)}{2!} \langle \phi^2 \rangle \right], \quad (20)$$

where, in the left hand of eq. (20), we have expanded to second order in ϕ the expectation value of $H^2(\varphi)$. Furthermore, in the right hand we have neglected the terms $\langle (\nabla \phi)^2 \rangle / (2a^2)$ and $\langle \dot{\phi}^2 \rangle / 2$ because they are very small with respect to the other on cosmological scales.

It is easy to show that the friction force generates entropy. It was calculated by Hosoya and Sakagami [for details in the calculation see [7]]

$$\dot{S}(t) = \frac{\eta_{eff}}{T} \left[\dot{\phi}_c \right]^2, \quad (21)$$

where $S(t)$ is the entropy density.

To summarize, equations (17), (18), (19) and (20) describe completely the dynamics of the background field ϕ_c on the thermal bath taking into account the temporal evolution of the temperature due to the expansion of the universe and the self-interaction of the inflaton field in an universe globally isotropic, homogeneous and spatially flat on cosmological scales which is described by a FRW metric.

III. AN EXAMPLE

If we assume an $\phi_c \rightarrow -\phi_c$ invariant potential (13), the Hubble parameter that describes the evolution of the universe can be obtained from the Friedmann equation (20)

$$H_c(\phi_c) = \sqrt{\frac{\pi G}{3}} \lambda \phi_c^2 + H_0, \quad (22)$$

where $H_0 = 2\sqrt{\frac{\pi G}{3}} \frac{m^2}{\lambda}$ is the Hubble parameter at the end of inflation.

In this case $V'_{eff}(\phi_c, T)$ can be written as

$$V'_{eff}(\phi_c, T) = \mathcal{M}^2(T) \phi_c + \frac{\lambda_{eff}^2(T)}{6} \phi_c^3, \quad (23)$$

where the effective mass and self-interaction parameters are given by

$$\mathcal{M}^2(T) = m^2 + \frac{\lambda^2 T^2}{24} \left(1 - \frac{4m}{\pi^2 T}\right) + \frac{3.2^6}{\pi} H_0 T, \quad (24)$$

$$\lambda_{eff}^2(T) = \lambda^2 + 2^7 \sqrt{\frac{3^3 G}{\pi}} \lambda T. \quad (25)$$

The third term in eq. (24) is related to the cosmological constant $H_0 = \sqrt{\frac{8\pi G}{3}} \Lambda$. This means that the cosmological constant should be $\Lambda = \frac{m^4}{2\lambda^2} \ll G^{-2}$. Furthermore, the second term in (25) is due to the thermal effects in the self-interaction of the inflaton field.

The temporal evolution of the inflaton field goes as t^{-1}

$$\phi_c(t) = \frac{\sqrt{6}}{\lambda t}. \quad (26)$$

Furthermore, in this case we obtain that the asymptotic value for the squared field fluctuations $\langle \phi^2(\vec{x}, t) \rangle|_{t \gg t_0}$, is

$$\langle \phi^2(\vec{x}, t) \rangle|_{t \gg t_0} \simeq \frac{T^2}{12} + \frac{3.2^7}{\pi} H_0 T, \quad (27)$$

where the last term is due to the interaction of the inflaton field with the thermal bath in an expanding universe and the first one takes into account this interaction without consider the expansion of the universe. To obtain the temporal evolution of the temperature we must to solve the eq. (15). The exact solution is very difficult to obtain, but the numerical calculation with $\lambda = 10^{-7}$ and $m = 10^{-6} G^{-1/2}$ is shown in the figure (1). With the parameter values here used we obtain $H_0 = 2.05 \cdot 10^{-5} G^{-1/2}$ and $\Lambda = 5 \cdot 10^{-11} G^{-2}$. The effective friction parameter is many orders of magnitude bigger than H_c during all the inflationary stage, so that the thermodynamical equilibrium is guaranteed. Furthermore, η_{eff} in eq. (18) approaches asymptotically to η during inflation, due to the last contribution in eq. (18) becomes negligible for late times.

Figure (1) shows the evolution of the temperature $T(x)$, where $x = \log_{10}(t)$. Note that the temperature increases with time until $t \simeq 10^{4.1} G^{1/2}$, and it is sufficiently large to be possible baryogenesis. After it, the temperature decreases slowly to remain almost constant always up to $T \simeq 5 \cdot 10^{-6} G^{-1/2}$. Hence, at the end of inflation the temperature remains with values bigger than the inflaton mass ($T(x)/m > 5$), so that the high temperature limit used in the calculation is always valid. Furthermore, as can be shown the change of entropy (21) is always positive [see figure (2)]. In the example here studied $\dot{S}$ reach its maximum value for times closed to $t \simeq 10^{4.01} G^{1/2}$ to decrease monotonically at the end of inflation.

At the end of inflation the fluctuations (27) are dominated by the last term: $\langle \phi^2 \rangle|_e \simeq \frac{3.2^7}{\pi} H_0 T \simeq 1.254 \cdot 10^{-8} G^{-1}$. They are shown in the figure (3) and we can see that remains almost constant at the end of inflation. The important fact here is that thermal fluctuations are more important than quantum fluctuations $\langle \phi^2 \rangle_{quant} \simeq H_c^2/(4\pi)$ in spite of the fact that $H_c < T$ during the last stages of inflation in which the high temperature limit is valid.

On the other hand, the number of e-folds $N = \int_{t_0}^t H_c(t) dt$ is sufficiently large to assure the solution to the horizon/flatness problems: $N[t > 10^{6.5} G^{1/2}] > 60$.

IV. FINAL COMMENTS

In this paper we have used a consistent formalism to find the effective equation of motion for the thermodynamical average of the quantum scalar field φ . This equation agree with the obtained by Hosoya and Sakagami, who obtained an additional term justified by the change of temperature during inflation ($\dot{T} \neq 0$). Such that a variation of temperature introduces a new term in the friction parameter. However, as inflation evolves the additional term in η_{eff} becomes negligible and $\eta_{eff} \simeq \eta$ at the end of inflation.

Furthermore, it is very interesting to note that the effective mass of the inflaton field now has two additional terms; one is product of the interaction [the second one in eq. (24)] and the other is originated by the cosmological constant

and the thermal effects [the third one in eq. (24)]. On the other hand, the thermal effects are also responsible for an additional term in the effective self-interaction parameter λ_{eff} [see eq. (25)]. Notice that $\lambda_{eff} \rightarrow 0$ as $\lambda \rightarrow 0$.

To summarize, in this paper I demonstrated that thermal effects originated by dissipation play an important role during inflation and do not must to be ignored in models for the early universe. Dissipation is so important that there are no oscillations of the inflaton field when it reach the minimum energetic configuration and at the same time, provides a smooth transition to the radiation era.

ACKNOWLEDGEMENTS

I would like to acknowledge CONICET (Argentina) for partial support and Universidad Nacional de Mar del Plata for financial support in the form of a research grant.

-
- [1] A. H. Guth, Phys. Rev. **D23**, 347 (1981).
 - [2] A. H. Guth and E. J. Weinberg, Nucl. Phys. **B212**, 321 (1983).
 - [3] A. D. Linde, Rep. Prog. Phys. **47**, 925 (1984); P. J. Steinhardt, Comments Nucl. Part. Phys. **12**, 273 (1984).
 - [4] A. D. Linde, Phys. Lett. **B108**, 389 (1982).
 - [5] A.D.Linde, *Particle Physics and Inflationary Cosmology* (Harwood, Chur, Switzerland, 1990) and references therein.
 - [6] A.A.Starobinsky, in *Current Topics in Field Theory, Quantum Gravity, and Strings*, ed. by H.J. de Vega and N.Sánchez, Lecture Notes in Physics 226 (Springer, New York, 1986).
 - [7] A. Hosoya and M. Sakagami, Phys. Rev. **D29**, 2228 (1984).
 - [8] M. Morikawa, Phys. Rev. **D33**, 3607 (1986).
 - [9] I. D. Lawrie, Phys. Rev. **D40**, 3330 (1989).
 - [10] M. Gleiser and R. O. Ramos, Phys. Rev. **D50**, 2441 (1994).
 - [11] A. Berera, Phys. Rev. Lett. **75**, 3218 (1995); Phys. Rev. **D54**, 2519 (1996); H. P. de Oliveira and R. O. Ramos, Phys. Rev. **D57**, 741 (1998); M. Bellini, Phys. Lett. **B428**, 31 (1998); Phys. Rev. **D58**, 103518 (1998)[Erratum-ibid. **D60**, 089901 (1999)]; J. M. F. Maia and J. A. S. Lima, Phys. Rev. **D60**, 101301 (1999).
 - [12] A. Berera, M. Gleiser and R. O. Ramos, Phys. Rev. **D58**, 123508 (1998).
 - [13] A. Berera, M. Gleiser and R. O. Ramos, Phys. Rev. Lett. **83**, 264 (1999).
 - [14] M. Bellini, Phys. Rev.**D63**, 123510 (2001); Gen. Rel. Grav. **35**, 41 (2003); Phys. Rev. **D64**, 123508 (2001); Phys. Rev. **D67**, 027303 (2003).

FIG. 1. Evolution of temperature $T(x)$. The time is shown in scale $x = \log_{10}(t)$.

FIG. 2. Evolution of change of entropy $\dot{S}(x)$. The time is shown in scale $x = \log_{10}(t)$.

FIG. 3. Evolution of fluctuations $\langle \phi^2 \rangle^{1/2}$. The time is shown in scale $x = \log_{10}(t)$.